

Revised, September 1998

HU Berlin-EP-98/40

hep-th@xxx/9807093

$Q\bar{Q}$ potential from AdS-CFT relation at $T \geq 0$: Dependence on orientation in internal space and higher curvature corrections

H. Dorn, H.-J. Otto ¹

*Humboldt-Universität zu Berlin
Institut für Physik, Theorie der Elementarteilchen
Invalidenstraße 110, D-10115 Berlin*

Abstract

Within the classical approximation we calculate the static $Q\bar{Q}$ potential via the AdS/CFT relation for nonzero temperature and arbitrary internal orientation of the quarks. We use a higher order curvature corrected target space background. For timelike Wilson loops there arises a critical line in the orientation-distance plane which is shifted to larger distances relative to the calculation with uncorrected background. Beyond that line there is no $Q\bar{Q}$ -force. The overall vanishing of the force for antipodal orientation known from zero temperature remains valid. The spacelike Wilson loops yield a string tension for a (2+1)-dimensional gauge theory, independent of the relative internal orientation, but sensitive to the background correction.

¹e-mail: dorn@physik.hu-berlin.de otto@physik.hu-berlin.de

1 Introduction

Part of the recent interest in relations between gauge field theories and string theory/supergravity on certain nontrivial backgrounds [1, 3, 2] is due to a conjecture expressing the Wilson loops in the gauge theory by the partition function of a string fulfilling boundary conditions set by the geometry of the Wilson loop [4, 5, 6]. This has been used to calculate the heavy quark-antiquark potential for $\mathcal{N} = 4$ super Yang-Mills in 3+1 dimensions in the t'Hooft limit and large effective coupling from the classical string action in $\text{AdS}_5 \times S^5$ [4, 5]. The procedure has been extended to general dimensions and to geometries arising as some limit of near extremal D-brane configurations [4, 6, 7, 8]. These background configurations with compactified Euclidean time coordinate are related to equilibrium $T > 0$ field theory. Since the periodicity condition breaks supersymmetry one gains access to the quark-antiquark potential in non-supersymmetric gauge theories (hopefully QCD) in one dimension lower [6, 7, 8].

The present paper will be concerned mainly with two aspects of this kind of calculations. In the $\text{AdS}_5 \times S^5$ calculation of ref.[4] a Coulombic potential arises, which is switched off for quark-antiquarks oriented antipodally with respect to the internal S^5 . This is in agreement with an argument telling that the corresponding configuration is of BPS type [9, 4]. We would like to know what happens in this antipodal case for $T > 0$, where supersymmetry is broken.

The classical approximation used on the string side of the duality is justified in the t'Hooft limit for large effective coupling, corresponding to small curvature of $\text{AdS}_5 \times S^5$ or of the relevant region of the near extremal D-brane configuration for $T > 0$. A complete treatment has to include both quantum corrections and corrections of the target space background, which after all has to be a solution for the effective string action exact in α' . As a first step in this direction we perform the classical calculation using the next order in α' corrected background of ref.[11, 12]. At $T > 0$ the quark-antiquark potential vanishes beyond some critical distance [7, 8]. This total screening is expected to be weakened if the corrections just mentioned are taken into account [10]. Therefore, our focus will be on the behaviour of the critical line in the $(\Delta\Theta, L)$ -plane (relative S^5 orientation and distance), if the correction of [11, 12] is switched on.

Our 10D metric for non-zero temperature $T > 0$ is given by

$$ds^2 = \alpha' \left\{ \frac{U^2}{R^2} \left(1 - \frac{U_T^4}{U^4}\right) e^{\gamma A(\frac{U_T}{U})} (dx^0)^2 + \frac{U^2}{R^2} e^{\gamma C(\frac{U_T}{U})} (dx^i)^2 \right. \\ \left. + \frac{R^2}{U^2 \left(1 - \frac{U_T^4}{U^4}\right)} e^{\gamma B(\frac{U_T}{U})} (dU)^2 + R^2 e^{\gamma D(\frac{U_T}{U})} (d\Omega_5)^2 \right\} = G_{MN} dx^M dx^N, \quad (1)$$

with ²

$$U_T = \pi R^2 T (1 + 15\gamma)^{-1}, \quad (2)$$

²The dimensionless parameter $\frac{1}{R}$ controls the curvature and is related to Yang-Mills quantities via $R^2 = \sqrt{2g_{YM}^2 N}$, which is equal to the effective coupling in the t'Hooft limit.

$$\begin{aligned}
A(\rho) &= -75\rho^4 - \frac{1225}{16}\rho^8 + \frac{695}{16}\rho^{12}, \\
B(\rho) &= 75\rho^4 + \frac{1175}{16}\rho^8 - \frac{4585}{16}\rho^{12}, \\
C(\rho) &= -\frac{25}{16}\rho^8 - \frac{25}{16}\rho^{12}, \\
D(\rho) &= \frac{15}{16}\rho^8 + \frac{15}{16}\rho^{12}.
\end{aligned} \tag{3}$$

We consider

$$\gamma = 0 \quad \text{or} \quad \gamma = \frac{1}{8}\zeta(3) R^{-6} \tag{4}$$

to describe either the near extremal D-3 brane metric used in refs. [7, 8] or the corresponding first order higher curvature corrected metric studied in ref. [11, 12]. In these target space metrics we calculate the Nambu-Goto action

$$S = \frac{1}{2\pi\alpha'} \int d\tau d\sigma \sqrt{\det(G_{MN} \partial_\mu x^M \partial_\nu x^N)} \tag{5}$$

for stationary ($\delta S = 0$) string world surfaces fulfilling certain boundary conditions at $U \rightarrow \infty$. In a first case, called timelike, the boundary of the string world sheet at $U \rightarrow \infty$ consists out of two lines running at ³

$$x^1 = \pm \frac{L}{2}, \quad x^2 = x^3 = 0, \quad \vec{\Theta} = \vec{\Theta}_Q \quad \text{or} \quad \vec{\Theta}_{\bar{Q}} \tag{6}$$

parallel to the x^0 -axis one times around the compactified x^0 , with compactification length $1/T$. In the second case, called spacelike, the boundary lines at $U \rightarrow \infty$ run parallel to the x^2 -axis at

$$x^1 = \pm \frac{L}{2}, \quad x^0 = x^3 = 0, \quad \vec{\Theta} = \vec{\Theta}_Q \quad \text{or} \quad \vec{\Theta}_{\bar{Q}}. \tag{7}$$

Via the CFT/AdS conjecture specified for Wilson loops in [4, 5, 6] one has in classical approximation for case 1

$$S_{|\text{case 1}} = \frac{1}{T} \cdot V_1(L, \Delta\Theta, T) \tag{8}$$

and for case 2 (Y measures the large x^2 interval.)

$$S_{|\text{case 2}} = Y \cdot V_2(L, \Delta\Theta), \tag{9}$$

where $V_1(L, \Delta\Theta, T)$ is the static quark-antiquark potential in 3-dimensional space at temperature T for separation L and an angle $\Delta\Theta$ between the internal orientation $\vec{\Theta}_Q$ and $\vec{\Theta}_{\bar{Q}}$ in S_5 . $V_2(L, \Delta\Theta)$ for $Y, L \gg \frac{1}{T}$ is the potential in 3-dimensional space-time (i.e. 2-dim. space) at zero temperature [6].

³ $\vec{\Theta}$ with $|\vec{\Theta}| = 1$ parametrises S^5 .

2 Timelike Wilson lines

In this section it is convenient to choose the gauge

$$\tau = x^0, \quad \sigma = x^1. \quad (10)$$

Furthermore, due to the symmetry of the problem, it is sufficient to look for world surfaces with $\partial_\tau x^M = \delta_0^M$, $\partial_\sigma x^2 = \partial_\sigma x^3 = 0$, only. Then the action becomes (' denotes derivatives with respect to σ , Θ is an angle variable on the great circle on S^5 passing $\vec{\Theta}_Q$ and $\vec{\Theta}_{\bar{Q}}$.)

$$S = \frac{1}{2\pi T} \int_{-\frac{L}{2}}^{+\frac{L}{2}} d\sigma e^{\frac{\gamma}{2}A(\frac{U_T}{\sigma})} \sqrt{\frac{U^4 - U_T^4}{R^4} e^{\gamma C(\frac{U_T}{\sigma})} + U'^2 e^{\gamma B(\frac{U_T}{\sigma})} + \Theta'^2 \frac{U^4 - U_T^4}{U^2} e^{\gamma D(\frac{U_T}{\sigma})}}. \quad (11)$$

The wanted solutions $U(\sigma)$, $\Theta(\sigma)$ are most easily found by making use of the conservation laws derived from the Lagrangian $\mathcal{L}$ ($S = \frac{1}{2\pi T} \int d\sigma \mathcal{L}$)

$$\begin{aligned} C_1 &= \frac{\partial \mathcal{L}}{\partial U'} U' + \frac{\partial \mathcal{L}}{\partial \Theta'} \Theta' - \mathcal{L}, \\ C_2 &= \frac{\partial \mathcal{L}}{\partial \Theta'}. \end{aligned} \quad (12)$$

To be as close as possible to ref. [4] we parametrise the conserved quantities C_1 and C_2 by $U_0 = U(0)$ and a parameter l via ⁴

$$\begin{aligned} C_2 &= U_0 l, \\ C_1 &= -\frac{U_0^2}{R^2} e^{\frac{\gamma}{2}\{C(\delta)-D(\delta)\}} \sqrt{(1-\delta^4)e^{\gamma\{A(\delta)+D(\delta)\}} - l^2}, \end{aligned} \quad (13)$$

with

$$\delta = \frac{U_T}{U_0}. \quad (14)$$

Then using the notation

$$f_\gamma(y, \delta) = (y^4 - \delta^4) e^{\gamma\{A(\frac{\delta}{y})+C(\frac{\delta}{y})-C(\delta)+D(\delta)\}} - l^2 y^2 e^{\gamma\{C(\frac{\delta}{y})-D(\frac{\delta}{y})+D(\delta)-C(\delta)\}} \quad (15)$$

straightforward integration yields

$$\sigma(U) = \frac{R^2}{U_T} \delta \sqrt{f_\gamma(1, \delta)} \int_1^{\frac{U}{U_0}} \frac{e^{\frac{\gamma}{2}\{B(\frac{\delta}{y})-C(\frac{\delta}{y})\}} dy}{\sqrt{y^4 - \delta^4} \sqrt{f_\gamma(y, \delta) - f_\gamma(1, \delta)}}, \quad (16)$$

$$\Theta(U) = l e^{\frac{\gamma}{2}\{D(\delta)-C(\delta)\}} \int_1^{\frac{U}{U_0}} \frac{y^2 e^{\frac{\gamma}{2}\{B(\frac{\delta}{y})+C(\frac{\delta}{y})-2D(\frac{\delta}{y})\}} dy}{\sqrt{y^4 - \delta^4} \sqrt{f_\gamma(y, \delta) - f_\gamma(1, \delta)}}, \quad (17)$$

⁴Making use of $U'(0) = 0$.

and in particular, using (2), for $L = 2\sigma(U \rightarrow \infty)$, $\Delta\Theta = 2\Theta(U \rightarrow \infty)$

$$L \cdot T = \frac{2}{\pi} (1 + 15\gamma)\delta \sqrt{f_\gamma(1, \delta)} \int_1^\infty \frac{e^{\frac{\gamma}{2}\{B(\frac{\delta}{y}) - C(\frac{\delta}{y})\}} dy}{\sqrt{y^4 - \delta^4} \sqrt{f_\gamma(y, \delta) - f_\gamma(1, \delta)}} , \quad (18)$$

$$\Delta\Theta = 2l e^{\frac{\gamma}{2}\{D(\delta) - C(\delta)\}} \int_1^\infty \frac{y^2 e^{\frac{\gamma}{2}\{B(\frac{\delta}{y}) + C(\frac{\delta}{y}) - 2D(\frac{\delta}{y})\}} dy}{\sqrt{y^4 - \delta^4} \sqrt{f_\gamma(y, \delta) - f_\gamma(1, \delta)}} . \quad (19)$$

Inserting the solution into the action functional (11) and making use of the conservation laws (12),(13) again, we get via (8) the static $Q\bar{Q}$ potential. Due to the behaviour of the integrand at large U it is divergent. With a preliminary cutoff Λ it looks like

$$V_1^{(\Lambda)} = \frac{U_T}{\pi} \delta^{-1} \int_1^{\frac{\Lambda}{U_0}} \frac{\sqrt{y^4 - \delta^4} e^{\frac{\gamma}{2}\{2A(\frac{\delta}{y}) + B(\frac{\delta}{y}) + C(\frac{\delta}{y}) + D(\delta) - C(\delta)\}} dy}{\sqrt{f_\gamma(y, \delta) - f_\gamma(1, \delta)}} . \quad (20)$$

Defining a renormalised potential along the lines of [4, 7, 8] requires a comment. The potential $V_1^{(\Lambda)}$ is linearly divergent. Subtracting the divergence yields a renormalised V_1 , fixed up to a finite additive constant which per se is completely irrelevant. From the experience with the $\Delta\Theta = 0$, $\gamma = 0$ case [7, 8], we expect that beyond some critical distance the configuration of two strings stretching straightly from $U = \Lambda$ to the horizon at $U = U_T$ at fixed values $x_1 = \pm \frac{L}{2}$, $\Theta = \pm \frac{\Delta\Theta}{2}$ is energetically favoured in comparison to our smooth U-shaped configuration (16),(17). Choosing $2 \cdot \frac{\Lambda - U_T}{2\pi}$, i.e. twice the energy stored in a string stretching between U_T and Λ as the subtraction term, the competing piecewise straight string configuration has been normalised to zero in [7, 8]. As a consequence, in the L -region with $V_1 \geq 0$ the $Q\bar{Q}$ force vanishes.

To extend this framework to our modified background (1), we first of all have to check, whether the competing piecewise straight string configuration (pssc) ⁵ still has an action (5) independent of L . Since also with (1) we have $G_{00} = 0$ at $U = U_T$, the determinant of the induced metric vanishes for the part of the pssc extending along the horizon. Only the pieces of the pssc running off the horizon contribute:

$$V_{\text{pssc}}^{(\Lambda)} = \frac{U_T}{\pi} \int_1^{\frac{\Lambda}{U_T}} e^{\frac{\gamma}{2}\{A(y^{-1}) + B(y^{-1})\}} dy . \quad (21)$$

For $\gamma = 0$ this reduces to $\frac{\Lambda - U_T}{\pi}$.

Now we choose $V_{\text{pssc}}^{(\Lambda)}$ as the subtraction constant to get a renormalised V_1 . Then the potential of the competing pssc is zero again, and our U-shaped configuration has a potential

$$V_1 = \lim_{\Lambda \rightarrow \infty} (V_1^{(\Lambda)} - V_{\text{pssc}}^{(\Lambda)}) ,$$

⁵At $x^1 = -\frac{L}{2}$, $\Theta = -\frac{\Delta\Theta}{2}$ from $U = \Lambda$ to $U = U_T$, then along the horizon to $x^1 = +\frac{L}{2}$, $\Theta = +\frac{\Delta\Theta}{2}$, and then at fixed x^1 , Θ back to $U = \Lambda$.

leading with (2) to

$$\begin{aligned} \frac{V_1}{R^2 T} = & (1 + 15\gamma)^{-1} \left\{ \delta^{-1} \int_1^\infty \left(\frac{\sqrt{y^4 - \delta^4} e^{\frac{\gamma}{2}\{2A(\frac{\delta}{y}) + B(\frac{\delta}{y}) + C(\frac{\delta}{y}) + D(\delta) - C(\delta)\}}}{\sqrt{f_\gamma(y, \delta) - f_\gamma(1, \delta)}} - 1 \right) dy \right. \\ & \left. + 1 - \delta^{-1} + \int_1^\infty \left(1 - e^{\frac{\gamma}{2}\{A(y^{-1}) + B(y^{-1})\}} \right) dy \right\}. \end{aligned} \quad (22)$$

The potential V_1 as a function of L and $\Delta\Theta$ is obtained by inserting in (22) the inversion of (18),(19), i.e. $\delta = \delta(L, \Delta\Theta)$ and $l = l(L, \Delta\Theta)$. The qualitative features of this inversion problem are best understood by looking at the contour lines of L , $\Delta\Theta$ and V_1 in the (l^2, δ^4) -plane. Since we are discussing string configurations with $U_T \leq U_0 < \infty$ we have $0 \leq \delta \leq 1$. In addition, the $\Delta\Theta$ -formula requires $l^2 \geq 0$, while the L -formula needs $l^2 \leq (1 - \delta^4)e^{\gamma\{A(\delta) + D(\delta)\}}$. Altogether our set of formulae is applicable in the region

$$\mathcal{G} = \{ (l^2, \delta^4) \mid 0 \leq l^2 \leq (1 - \delta^4)e^{\gamma\{A(\delta) + D(\delta)\}}, \delta^4 \geq 0 \}. \quad (23)$$

A generic feature of these integrals is their monotonous increase with l^2 at fixed δ . As a consequence, V_1 and $\Delta\Theta$ are monotonously increasing functions of l^2 . Due to the inverse behaviour of the square root prefactor it is difficult to make a definite statement concerning L .

The following discussion of the contour line maps is based on explicit analysis of the $\gamma = 0$ case. However, the qualitative features, forced by the singularities of the functions under discussion, are not changed by switching on γ . We start with the $\Delta\Theta$ lines in $0 \leq \delta^4 \leq 1$, $0 \leq l^2 \leq 2$. Here $(l^2, \delta^4) = (0, 1)$ is a singular point⁷. Across this point run all contour lines from $\Delta\Theta = 0$ up to $\Delta\Theta = \infty$. The $\Delta\Theta = 0$ line goes straight down to the point $(0, 0)$, the $\Delta\Theta = \infty$ line first runs parallel to the l^2 axis to $(2, 1)$ and then straight down to $(2, 0)$. The $\Delta\Theta = \pi$ line goes through $(l^2, \delta^4) = (1, 0)$.

Let us turn to the L -lines. Again $(0, 1)$ is a singular point. Here the $L = \infty$ line at $\delta^4 = 1$, $-\infty < l^2 < 0$ meets the straight zero-line connecting $(1, 0)$ with $(0, 1)$. The approach of all lines with $0 \leq L < \infty$ to the singular point is possible due to the balance between the logarithmic divergent integral for $\delta^4 \rightarrow 1$ and the vanishing square root prefactor. Therefore, *all* lines with $L > 0$ start tangent to the $L = \infty$ line into the region $l^2 < 0$. Those with small enough L must turn to the $l^2 > 0$ region, since they must stay close to the zero line also away from the singular point. Altogether L -lines up to some maximal value $L_{\max}$ cross the δ^4 -axis twice. For $l^2 = 0$ the inversion of (18) exists only for $L \leq L_{\max}$, for the $\gamma = 0$ case see also [7, 8]. From the point where the $L_{\max}$ -line touches the δ^4 -axis to the point $(l^2, \delta^4) = (1, 0)$ extends a ridge of decreasing height. For a given $0 \leq l^2 < 1$ the inverse of (18) exists for values of L smaller than the corresponding ridge height, only.

⁶Using the fact that $A(\rho) + D(\rho)$ as well as $C(\rho) - D(\rho)$ in $0 \leq \rho \leq 1$ are negative and monotonously decreasing one can convince oneself that the integrals in (18),(19) and (22) are well defined *inside* $\mathcal{G}$ for arbitrary γ . They exist even in a larger region, for the simplest case $\gamma = 0$ in $0 \leq \delta < 1$, $-\infty < l^2 < 2$.

⁷ Vanishing of the factor l versus logarithmic divergence of the integral for $\delta^4 \rightarrow 1$.




Fig.1 Critical lines $V_1 = 0$ as functions of $\frac{1}{\pi} \cdot \Delta\Theta$ (horizontal) and $\frac{\pi}{2}T \cdot L$ (vertical). The lowest line is for $\gamma = 0$, the others for $\gamma = 0.01, 0.02$, and 0.05 respectively.

Finally, the V_1 map is governed by a singularity with crossing contour lines at $(l^2, \delta^4) = (1, 0)$. On the l^2 -axis V_1 tends to $\infty(-\infty)$ for $l^2 > 1(< 1)$. Also $l^2 = 2$, $0 \leq \delta^4 < 1$ is a line of $V_1 = \infty$. On the contrary, V_1 takes finite values on $\delta^4 = 1$, $l^2 < 2$. In particular one finds $V_1 = 0$ at $(0, 1)$. The zero line of V_1 from $(0, 1)$ to $(1, 0)$ is the critical line we are after. Beyond this critical line there is no longer any force between the $Q\bar{Q}$ -pair. From the analysis of [7, 8] and a theorem in [13] we know that, for $\gamma = 0$, the zero line coming from $(0, 1)$ at first must go to the $l^2 < 0$ region to cross the δ^4 -axis at a value below the L -ridge. We have no general proof that the $V_1 = 0$ line is below the L -ridge in whole $\mathcal{G}$, but there is numerical evidence for this scenario for not too large γ .

After this qualitative discussion of the behaviour of the potential we present the results of a numerical evaluation. In fig.1 the critical lines, ($V_1 = 0$), in the $(\frac{1}{\pi} \cdot \Delta\Theta, \frac{\pi}{2}T \cdot L)$ -plane, obtained for various values of γ are shown. Beyond the critical line there is no $Q\bar{Q}$ force. Although for $\gamma = 0$, i.e. the zeroth order metric used in [7, 8], the potential is R -dependent, the $V_1 = 0$ contour line in the $(\Delta\Theta, L)$ -plane is independent of R . With the first order corrected metric of [11, 12] the R -dependence originates from the γ -dependence via (4). Obviously, the correction acts in the right direction, the critical line for the onset of total screening is shifted to larger distances over the whole range of $\Delta\Theta$. The chosen values of $\gamma = 0.01, 0.02$ and 0.05 correspond to $R^2 = 2.468, 1.959$ and 1.443 respectively.




Fig.2 $\frac{1}{R^2 T} \cdot V_1$ as a function of $\frac{\pi}{2} T \cdot L$ at $\gamma = 0$ for various values of $\frac{1}{\pi} \Delta \Theta$, from left to right: 0.8, 0.5, 0.

To get an impression of the influence of $\Delta \Theta$ on the potential we show in fig.2 $\frac{1}{R^2 T} \cdot V_1$ as a function of $\frac{\pi}{2} T \cdot L$ for various values of $\frac{1}{\pi} \cdot \Delta \Theta$ in the case of zeroth order metrics, i.e. $\gamma = 0$. The $Q\bar{Q}$ force is zero for distances larger than the value of the zero of the drawn potential. For illustration we also followed the potential a little bit for positive values. The cusp with the accompanying backward movement indicates that one goes beyond the L -ridge discussed above.

For a better understanding of the mechanism switching off the potential completely for $\Delta \Theta \rightarrow \pi$ we show (for $\gamma = 0$) in fig.3 the (l^2, δ^4) -plane with the region $\mathcal{G}$, the contour line $V_1 = 0$ and contour lines of $\Delta \Theta$. Any $\Delta \Theta$ -line below π enters $\mathcal{G}$ and the region of nonvanishing force for small enough values of δ^4 . The $\Delta \Theta = \pi$ -line touches $\mathcal{G}$ at the point $(1, 0)$ only. This figure also illustrates some statements made in the qualitative discussion above.

Finally, fig.4 shows, again for $\gamma = 0$, some contour lines of $\frac{\pi}{2} T \cdot L$ in the (l^2, δ^4) -plane. The numerical curves illustrate the general analytic arguments given above. The L -ridge is visible clearly. The L -line touching the δ^4 axis corresponds to $L_{\max} = 0.435$.




Fig.3 (l^2, δ^4) plane with region $\mathcal{G}$ for $\gamma = 0$ (shadowed). Contour lines of $\frac{1}{\pi} \cdot \Delta\Theta$ are shown for 0.5, 0.8, 1, from left to right. The dashed line is the $V_1 = 0$ contour line.




Fig.4 (l^2, δ^4) plane with region $\mathcal{G}$ for $\gamma = 0$ (shadowed). Contour lines of $\frac{\pi}{2}T \cdot L$ are shown for 0.5, 0.435, 0.3, 0.2, 0.1, from left to right. The dashed line is the $V_1 = 0$ contour line.

Closing this section we present the results of an expansion of (18),(19),(22) for small δ . It is related to either $T \rightarrow 0$ at fixed distance or to small distances at fixed temperature. Within this expansion one encounters the following l^2 -dependent integrals, expressible by the hypergeometric function F , the elliptic integrals K and E as ($n \geq 0$)

$$\begin{aligned}
b_{2n}(l) &= \int_1^\infty \frac{dy}{y^{2n} \sqrt{(y^2-1)(y^2+1-l^2)}} = \frac{1}{2} B(n + \frac{1}{2}, \frac{1}{2}) F(n + \frac{1}{2}, \frac{1}{2}, n+1, l^2-1) , \\
b_{-2}(l) &= \int_1^\infty \left(\frac{y^2}{\sqrt{(y^2-1)(y^2+1-l^2)}} - 1 \right) dy \\
&= 1 + \frac{\pi}{2\sqrt{2-l^2}} F(\frac{1}{2}, \frac{1}{2}, 1, \frac{1-l^2}{2-l^2}) - \frac{\pi}{2} \sqrt{2-l^2} F(-\frac{1}{2}, \frac{1}{2}, 1, \frac{1-l^2}{2-l^2}) . \tag{24}
\end{aligned}$$

All $b(l)$ are monotonously increasing with l^2 . Special values are

$$\begin{aligned}
b_{2n}(0) &= \frac{1}{4} B(\frac{n}{2} + \frac{1}{4}, \frac{1}{2}) , \\
b_{-2}(0) &= 1 + \frac{1}{\sqrt{2}} \left(K(\frac{\sqrt{2}}{2}) - 2E(\frac{\sqrt{2}}{2}) \right) \approx 0.401 , \\
b_{2n}(1) &= \frac{1}{2} B(n + \frac{1}{2}, \frac{1}{2}) , \\
b_{-2}(1) &= 1 . \tag{25}
\end{aligned}$$

Then we get ($c(\gamma) = \int_1^\infty (1 - e^{\frac{\gamma}{2}(A(\frac{1}{x})+B(\frac{1}{x}))} dx)$)

$$\frac{V_1}{R^2 T} = \frac{1}{1+15\gamma} \left(\frac{b_{-2}(l)-1}{\delta} + 1 + c(\gamma) - \frac{1+75\gamma}{2} b_2(l) \delta^3 + O(\delta^7) \right) , \tag{26}$$

$$\begin{aligned}
L \cdot T &= \frac{2}{\pi} (1+15\gamma) \sqrt{1-l^2 - (1+75\gamma)\delta^4 + O(\delta^8)} \\
&\quad \cdot \left(b_2(l) \delta + \frac{1+75\gamma}{2} b_6(l) \delta^5 + O(\delta^9) \right) , \tag{27}
\end{aligned}$$

$$\Delta\Theta = 2l \left(b_0(l) + \frac{1+75\gamma}{2} b_4(l) \delta^4 + O(\delta^8) \right) . \tag{28}$$

The insertion of inversed (27),(28) into (26) is straightforward and will not be discussed here.

3 Spacelike Wilson loops

To describe a string world sheet satisfying the boundary condition (7) it is convenient to choose the gauge

$$\tau = x^2, \quad \sigma = x^1 . \tag{29}$$

The symmetry of the problem allows to restrict ourselves to world surfaces with $\partial_\tau x^M = \delta_2^M$, $\partial_\sigma x^0 = \partial_\sigma x^3 = 0$. Then the action becomes, compare (9),

$$S_{\text{case 2}} = \frac{Y}{2\pi} \int_{-\frac{L}{2}}^{+\frac{L}{2}} d\sigma e^{\frac{\gamma}{2}C(\frac{U_T}{U})} \cdot \sqrt{\frac{U^4}{R^4} e^{\gamma C(\frac{U_T}{U})} + U'^2 \frac{U^4}{U^4 - U_T^4} e^{\gamma B(\frac{U_T}{U})} + \Theta'^2 U^2 e^{\gamma D(\frac{U_T}{U})}}. \quad (30)$$

The conserved quantities $\tilde{C}_1$ and $\tilde{C}_2$ defined analogously to (12) will be parametrised by

$$\begin{aligned} \tilde{C}_2 &= U_0 l \\ \tilde{C}_1 &= -\frac{U_0^2}{R^2} e^{\gamma C(\delta)} \sqrt{1 - l^2 e^{-\gamma\{C(\delta)+D(\delta)\}}} . \end{aligned} \quad (31)$$

Then we find with the notation

$$g_\gamma(y, \delta) = y^4 e^{2\gamma\{C(\frac{\delta}{y})-C(\delta)\}} - y^2 l^2 e^{\gamma\{C(\frac{\delta}{y})-D(\frac{\delta}{y})-2C(\delta)\}}, \quad (32)$$

repeating the steps in section 2,

$$L \cdot T = \frac{2}{\pi} (1 + 15\gamma) \sqrt{g_\gamma(1, \delta)} \delta \int_1^\infty \frac{e^{\frac{\gamma}{2}\{B(\frac{\delta}{y})-C(\frac{\delta}{y})\}} dy}{\sqrt{y^4 - \delta^4} \sqrt{g_\gamma(y, \delta) - g_\gamma(1, \delta)}}, \quad (33)$$

$$\Delta\Theta = 2l \int_1^\infty \frac{y^2 e^{\frac{\gamma}{2}\{B(\frac{\delta}{y})+C(\frac{\delta}{y})-2D(\frac{\delta}{y})-2C(\delta)\}} dy}{\sqrt{y^4 - \delta^4} \sqrt{g_\gamma(y, \delta) - g_\gamma(1, \delta)}}, \quad (34)$$

$$\frac{V_2}{R^2 T} = (1 + 15\gamma)^{-1} \left\{ \delta^{-1} \int_1^\infty \left(\frac{y^4 e^{\frac{\gamma}{2}\{B(\frac{\delta}{y})+3C(\frac{\delta}{y})-2C(\delta)\}}}{\sqrt{y^4 - \delta^4} \sqrt{g_\gamma(y, \delta) - g_\gamma(1, \delta)}} - 1 \right) + 1 - \delta^{-1} \right\}. \quad (35)$$

To define the renormalised potential V_2 we have chosen as the subtraction $\frac{\Lambda - U_T}{\pi}$. Contrary to the previous section the pssc configuration has a L -dependent action. This is due to different behaviour on the horizon, $G_{22} \neq 0$ versus $G_{00} = 0$. With the same subtraction as for V_2 one gets, e.g. for $\Delta\Theta = 0$,

$$V_{2,\text{pssc}} = \frac{U_T^2}{2\pi R^2} e^{\gamma C(1)} \cdot L + \frac{U_T}{\pi} \int_1^\infty \left(\frac{y^2 e^{\frac{\gamma}{2}\{B(\frac{1}{y})+C(\frac{1}{y})\}}}{\sqrt{y^4 - 1}} - 1 \right) dy. \quad (36)$$

We expect this to be energetically disfavoured over the whole range of L .

The $L - \delta$ relation differs qualitatively from that in section 2. Due to the absence of δ in the square root prefactor we can reach arbitrary large values of L by approaching the singularity of the integral at $\delta = 1$. Furthermore, L is monotonously increasing with δ^4 .

⁸ The leading terms in the expansion of all the integrals in (33)-(35) for $\delta \rightarrow 1$ coincide

⁸This is obviously for $\gamma = 0$. It can be demonstrated by simple estimates also for small enough values $\gamma > 0$.

due to the equality of the leading term of the expansion of the respective integrands for $y \rightarrow 1$. Denoting with a fixed $\epsilon > 0$

$$I(\epsilon, l) = \int_1^{1+\epsilon} \frac{dy}{\sqrt{y^4 - \delta^4} \sqrt{g_\gamma(y, \delta) - g_\gamma(1, \delta)}} ,$$

we get

$$\begin{aligned} L \cdot T &= \frac{2}{\pi} (1 + 15\gamma) \sqrt{g_\gamma(1, 1)} e^{\frac{\gamma}{2}\{B(1)-C(1)\}} (I(\epsilon, l) + O(1)) , \\ \Delta\Theta &= 2l e^{\frac{\gamma}{2}\{B(1)-C(1)-2D(1)\}} (I(\epsilon, l) + O(1)) , \\ \frac{V_2}{R^2 T} &= (1 + 15\gamma)^{-1} e^{\frac{\gamma}{2}\{B(1)+C(1)\}} (I(\epsilon, l) + O(1)) . \end{aligned} \quad (37)$$

Using the first line of (37) to eliminate I in the last line one arrives at

$$V_2 = \frac{\pi R^2 T^2}{2} (1 - \frac{265}{8}\gamma) \cdot \frac{L}{\sqrt{1 - l^2 e^{\frac{5}{4}\gamma}}} + O(1) . \quad (38)$$

At the very end we want to know the potential for fixed $\Delta\Theta$. According to the second line of (37) at $L \rightarrow \infty$ this requires $l \rightarrow 0$. Using the first line of (37) to eliminate I in the second line one gets $l = \frac{\Delta\Theta}{\pi L T} e^{\gamma(15+D(1))} + o(\frac{1}{L})$ and altogether up to linear terms in γ

$$V_2(\Delta\Theta, L) = \frac{\pi R^2 T^2}{2} (1 - \frac{265}{8}\gamma) \cdot L + \frac{1}{4\pi} R^2 (\Delta\Theta)^2 (1 + \frac{15}{8}\gamma) \cdot \frac{1}{L} + O(1) . \quad (39)$$

The terms which we neglected and indicated as $O(1)$ are, up to eq.(38), more precisely $\propto L e^{-\pi T L}$, as follows by repeating the estimates of ref.[14]. There the case $\Delta\Theta = 0$ was studied and the absence of any Lüscher term $\propto 1/L$ in the large L expansion was pointed out to be an obstacle for the potential, derived in the approach of [4, 5, 6], to be identified with a QCD potential. We find it interesting that for $\Delta\Theta \neq 0$ an $1/L$ term arises. However, due to the wrong sign and a coupling dependence⁹ it cannot be interpreted as a Lüscher term. On the other side one should keep in mind that this term is also not Coulombic since in 2+1 dimensions the Coulomb term is $\propto \log L$.

4 Conclusions

We showed that within the classical approximation the static quark-antiquark potential is completely switched off for antipodal $Q\bar{Q}$ orientation in the internal S^5 also for $T > 0$. In contrast to the $T = 0$ case this is, due to the broken SUSY, *not* required by any BPS argument. The effect is present both for the zeroth order background metrics of refs.[7, 8] as well as for the higher curvature corrected background of ref.[11, 12]. One should add, that certainly this is an effect of the backgrounds used. In other circumstances, with BPS

⁹We thank P. Olesen for a comment on this fact.

arguments set out of work, nonvanishing $Q\bar{Q}$ forces may be possible for all orientations in internal space. Here, for example, we have in mind a $\mathcal{N} = 1$, $T = 0$ case based on a simple orbifold construction over S^5 [15].

The replacement of the lowest order background by the corrected ones shifts the critical line in the orientation-distance plane to larger distances for all values of the relative orientation in internal space. This shift acts in the right direction, if one follows the arguments of ref.[10] stating the vanishing of the total screening as soon as all corrections are taken into account.

From the study of spacelike Wilson loops we got the string tension for a temperature zero gauge theory in (2+1) dimensions. This tension is independent of the relative internal orientation of the quarks. However, the tension is sensible to switching on the string target space background corrections. The subleading terms in a large distance expansion of the potential are dependent on the relative internal orientation. In particular we found a nextleading $1/L$ term, absent for parallel orientation.

In this picture T plays the role of a regularization parameter which, to make contact with (2+1)-dimensional non-supersymmetric QCD, has to be sent to infinity. To keep a finite tension, R^2 has to go to zero. Obviously, our approximation is not suited to study this limit. However, our formula (39) suggests that the $\Delta\Theta$ dependent $1/L$ term drops out in this limit. This indeed should happen since there is no place in QCD for the internal S^5 orientation parameter which is a remnant of $\mathcal{N} = 4$ SUSY.

Acknowledgement: We would like to thank A. Karch, V. Pershin, Chr. Preitschopf and S. Theisen for useful discussions.

References

- [1] J. Maldacena, *The large N limit of superconformal field theories and supergravity*, hep-th/9711200
- [2] S.S. Gubser, I.R. Klebanov, A.M. Polyakov, *Gauge theory correlators from noncritical string theory*, hep-th/9802109
- [3] E. Witten, *Anti-De Sitter space and holography*, hep-th/9802150
- [4] J. Maldacena, *Phys. Rev. Lett.* **80** (1998) 4859, hep-th/9803002
- [5] S.-J. Rey, J.-T. Yee, *Macroscopic strings as heavy quarks in large N gauge theory and anti-de Sitter supergravity*, hep-th/9803001
- [6] E. Witten, *Anti-de Sitter space, thermal phase transition, and confinement in gauge theories*, hep-th/9803131
- [7] S.-J. Rey, S. Theisen, J.-T. Yee, *Wilson-Polyakov loop at finite temperature in large N gauge theory and anti-de Sitter supergravity*, hep-th/9803135

- [8] A. Brandhuber, N. Itzhaki, J. Sonnenschein, S. Yankielowicz, *Wilson loops in the large N limit at finite temperature*, hep-th/9803137
A. Brandhuber, N. Itzhaki, J. Sonnenschein, S. Yankielowicz, JHEP 06(1998)001, hep-th/9803263
- [9] C.G. Callan, J.M. Maldacena, *Nucl. Phys.* **513** (1998) 198, hep-th/9708147
- [10] D.J. Gross, H. Ooguri, *Aspects of large gauge theory dynamics as seen by string theory*, hep-th/9805129
- [11] S.S. Gubser, I.R. Klebanov, A.A. Tseytlin, *Coupling constant dependence in the thermodynamics of $N=4$ supersymmetric Yang-Mills theory*, hep-th/9805156
- [12] J. Pawelczyk, S. Theisen, *$AdS_5 \times S^5$ black hole metrics at $O((\alpha')^3)$* , hep-th/9808126
- [13] J.A. Minahan, N.P. Warner, *Quark potentials in the Higgs phase of large N supersymmetric Yang-Mills theories*, hep-th/9805104
- [14] J. Greensite, P. Olesen, *Remarks on the heavy quark potential in the supergravity approach*, hep-th/9806235
- [15] S. Kachru, E. Silverstein, *Phys. Rev. Lett.* **80** (1998) 4855, hep-th/9802183